

“In our experiment we see that young men who take ibuprofen resemble what we see in 70-year-old men. We don’t yet know what the consequences could be, and whether the conditions persists after you stop taking ibuprofen. Subsequent studies will look into this,” he says.

Read More: [Painkiller use during pregnancy damages baby boys' testicle development](#)[8]

Testes produce less testosterone

As part of the study, Kristensen’s colleagues analysed testicular tissue to study how the testes responded to ibuprofen. To do this, they studied tissue from coma patients who had a complete loss of brain function.

These laboratory experiments showed that the testes precipitation of testosterone was reduced when men received ibuprofen.

But it was not just testosterone: All hormones produced in the testes were affected by ibuprofen.

Read More: [Marriage reduces testosterone in men](#)[9]

First study of its kind

Professor Cecilia Ramlau-Hansen from the Department of Public Health at Aarhus University, Denmark, describes the study as “exciting” and “well executed.” She was not involved in the research.

“The scientists use a range of approaches to investigate how ibuprofen influences the balance of hormones and the results of their research support each other,” says Ramlau-Hansen.

One of the strengths of the study is the randomised trial, in which the male participants are split into two groups: One receiving ibuprofen, the other receiving a placebo. Randomised trials normally lead to good results, she says.

“As far as I know it’s the first time anyone has made a randomised control trial to test how ibuprofen effects hormone balance,” says Ramlau-Hansen.

Read More: [Infertility in men could point to more serious health problems later in life](#)[10]

Over-the-counter medicine can also have side effects

However, one drawback is that the study is not especially large with just 31 participants.

Moreover, the new study does not investigate how ibuprofen influences the hormone balance in the long term. They do not know whether the hormone imbalance returns to normal after taking ibuprofen for a certain period, or whether the hypogonadism becomes permanent.

The results indicate that men should use ibuprofen sparingly, even if it is available to purchase over-the-counter and without prescription, say Ramlau-Hansen and Kristensen.

Some people need to take ibuprofen to treat pain, such as arthritis patients. But if you have a sports injury or headache, you should consider not taking ibuprofen.

“Research shows that many athletes, both professional and amateurs, take ibuprofen if they have an injury but wish to continue playing anyway,” says Kristensen.

But the new study indicates that it might be better to take a break until the injury is healed. When you exercise, it is especially important to have a balance of testosterone, which plays an important role in muscle development, he says.

Read More: [Researchers are looking for young impotent men](#)[11]

Previous studies reveal mixed results

Kristensen and colleagues previously showed that boys have an increased risk of birth defects of their sexual organs if their mother took paracetamol painkillers while pregnant.

Other research into painkillers and hormone imbalance produced mixed results, says Ramlau-Hansen.

Some studies have shown that painkillers, such as ibuprofen and paracetamol, can disturb the hormone balance and reduce the testes production of testosterone. Other studies have indicated that painkillers can reduce sperm quality.

Another large population study by Ramlau-Hansen did not find any connection between painkillers and a reduced testosterone level.

[Read more in the Danish version of this story on Videnskab.dk](#) [12]

 [Ibuprofen could reduce testosterone production in men. \(Photo: Wikipedia\)](#) [13]

 [Scientists do not know whether ibuprofen does any permanently damage to men's endocrine system. \(Photo: Shutterstock\)](#) [14]

 [Ibuprofen could reduce testosterone production in men. \(Photo: Wikipedia\)](#) [15]

[Common painkillers can cause blood clots](#) [16] [Decreasing painkiller use lowers symptoms of depression and anxiety](#) [17] [Few regular users of strong painkillers](#) [18] [Painkiller use during pregnancy damages baby boys' testicle development](#) [19] [Painkillers lower fertility in mice](#) [7]

[David Møbjerg Kristensen](#) [20] [Cecilia Ramlau-Hansen](#) [21]

['Ibuprofen alters human testicular physiology to produce a state of compensated hypogonadism', PNAS 2018.](#) [6]

[Anne Ringgaard](#) [22]

Catherine Jex

February 13, 2018 - 08:00

This field is not in use. The footer is displayed in the mini panel called "Footer (mini panel)"

Source URL: <http://sciencenordic.com/ibuprofen-can-damage-men%E2%80%99s-endocrine-system>

Links:

[1] <http://sciencenordic.com/category/section/health>

[2] <http://sciencenordic.com/hormones>

[3] <http://sciencenordic.com/painkillers>

[4] <http://sciencenordic.com/category/countries/denmark>

- [5] <http://sciencenordic.com/videnskabdk>
- [6] <http://www.pnas.org/content/early/2018/01/03/1715035115>
- [7] <http://sciencenordic.com/painkillers-lower-fertility-mice>
- [8] <http://sciencenordic.com/painkiller-use-during-pregnancy-damages-baby-boy?s-testicle-development>
- [9] <http://sciencenordic.com/marriage-reduces-testosterone-men>
- [10] <http://sciencenordic.com/infertility-men-could-point-more-serious-health-problems-later-life>
- [11] <http://sciencenordic.com/researchers-are-looking-young-impotent-men>
- [12] <https://videnskab.dk/krop-sundhed/ibuprofen-kan-skade-maends-hormonsystem>
- [13] <http://sciencenordic.com/sites/default/files/ibuprofen.jpg>
- [14] http://sciencenordic.com/sites/default/files/ung_mand_smertestillende.jpg
- [15] http://sciencenordic.com/sites/default/files/ibuprofen_0.jpg
- [16] <http://sciencenordic.com/common-painkillers-can-cause-blood-clots>
- [17] <http://sciencenordic.com/decreasing-painkiller-use-lowers-symptoms-depression-and-anxiety>
- [18] <http://sciencenordic.com/few-regular-users-strong-painkillers>
- [19] <http://sciencenordic.com/painkiller-use-during-pregnancy-damages-baby-boy%E2%80%99s-testicle-development>
- [20] [https://research.regionh.dk/rigshospitalet/da/persons/david-moebjerg-kristensen\(ba351c36-7044-4f60-ad0b-f42e48e8e3f3\).html](https://research.regionh.dk/rigshospitalet/da/persons/david-moebjerg-kristensen(ba351c36-7044-4f60-ad0b-f42e48e8e3f3).html)
- [21] <http://pure.au.dk/portal/en/persons/cecilia-ramlauhansen%2876e02a27-dc4a-4c88-968c-d82d47f9961f%29.html>
- [22] <http://sciencenordic.com/content/anne-ringgaard>